

Package ‘ptsuite’

April 2, 2019

Title Tail Index Estimation for Power Law Distributions

Version 1.0.0

Description Various estimation methods for the shape parameter of Pareto distributed data. This package contains functions for various estimation methods such as maximum likelihood (Newman, 2005)<doi:10.1016/j.cities.2012.03.001>, Hill's estimator (Hill, 1975)<doi:10.1214/aos/1176343247>, least squares (Zaher et al., 2014)<doi:10.9734/BJMCS/2014/10890>, method of moments (Rytgaard, 1990)<doi:10.2143/AST.20.2.2005443>, percentiles (Bhatti et al., 2018)<doi:10.1371/journal.pone.0196456>, and weighted least squares (Nair et al., 2019) to estimate the shape parameter of Pareto distributed data. It also provides both a heuristic method (Hubert et al., 2013)<doi:10.1016/j.csda.2012.07.011> and a goodness of fit test (Gulati and Shapiro, 2008)<doi:10.1007/978-0-8176-4619-6> for testing for Pareto data as well as a method for generating Pareto distributed data.

Depends R (>= 3.5.0)

License GPL-3

LazyData true

LinkingTo Rcpp

Imports Rcpp

RoxygenNote 6.1.1

Suggests plotly

NeedsCompilation yes

Author Ranjiva Munasinghe [aut],
Pathum Kossinna [cre, aut],
Dovini Jayasinghe [aut],
Dilanka Wijeratne [aut]

Maintainer Pathum Kossinna <pathum@mindlanka.org>

Repository CRAN

Date/Publication 2019-04-02 15:10:03 UTC

R topics documented:

ptsuite-package	2
alpha_geometric_percentile	3
alpha_hills	4
alpha_ls	5
alpha_mle	6
alpha_modified_percentile	7
alpha_moment	7
alpha_percentile	8
alpha_wls	9
generate_all_estimates	10
generate_pareto	10
pareto_qq_test	11
pareto_test	12

Index	13
-------	----

ptsuite-package *ptsuite: Package for Pareto Parameter Estimation*

Description

The ptsuite package provides functions to estimate parameters of Type I Pareto data. The details of the methods used and the equations implemented are given in the vignette. The package contains the following functions:

Estimator Functions

- Maximum Likelihood Estimator {[alpha_mle](#)}
- Weighted Least Squares Estimator {[alpha_wls](#)}
- Hill's Estimator {[alpha_hills](#)}
- Method of Moments Estimator {[alpha_moment](#)}
- Method of Percentiles Estimator {[alpha_percentile](#)}
- Method of Modified Percentiles Estimator {[alpha_modified_percentile](#)}
- Method of Geometric Percentiles {[alpha_geometric_percentile](#)}
- Least Squares Estimator {[alpha_ls](#)}

Other Functions

- Generate Pareto Data {[generate_pareto](#)}
- Estimates from all estimators {[generate_all_estimates](#)}
- Q-Q Plot to test for Pareto Distribution {[pareto_qq_test](#)}
- Pareto Test {[pareto_test](#)}

References

- Newman MEJ (2005). "Power Laws, Pareto Distributions And Zipf's Law." *Contemporary Physics*, 46, 323-351.
- Nair J, Wierman A, Zwart B (2019). "The Fundamentals Of Heavy Tails: Properties, Emergence, And Identification." <http://users.cms.caltech.edu/adamw/heavytails.html>.
- Pokorna M (2016). Estimation and Application of the Tail Index. Bachelor's thesis, Charles University in Prague, Faculty of Social Sciences, Institute of Economic Studies.
- Hill B (1975). "A Simple General Approach To Inference About The Tail Of A Distribution." *The Annals of Statistics*, 3(5), 1163-1174.
- Rytgaard M (1990). "Estimation In The Pareto Distribution." *ASTIN Bulletin: The Journal Of The IAA*, 20(2), 201-216.
- Brazauskas V, Serfling R (2000). "Robust and Efficient Estimation Of The Tail Index Of A Single-Parameter Pareto Distribution." *North American Actuarial Journal*, 4, 12-27.
- Bhatti SH, Hussain S, Ahmad T, Aslam M, Aftab M, Raza MA (2018). "Efficient estimation of Pareto model: Some modified percentile estimators." *PLoS ONE*, 13(5), 1-15.
- Zaher HM, El-Sheik AA, El-Magd NATA (2014). "Estimation of Pareto Parameters Using a Fuzzy Least-Squares Method and Other Known Techniques with a Comparison." *British Journal of Mathematics & Computer Science*, 4(14), 2067-2088.
- Gulati S, Shapiro S (2008). "Goodness-of-Fit Tests for Pareto Distribution." In F Vonta (ed.), *Statistical Models and Methods for Biomedical and Technical Systems*, chapter 19, pp. 259-274. Birkhauser Basel. ISBN 978-0-8176-4619-6. doi:10.1007/978-0-8176-4619-6.

Description

This function uses the Geometric Method of Percentiles to estimate the shape parameter of a given set of data. (Bhatti et al. 2018)

Usage

```
alpha_geometric_percentile(dat)
```

Arguments

dat	vector of observations
-----	------------------------

Value

A list of the following form:

shape Estimate of the shape parameter of the data

scale Estimate of the scale parameter of the data (which is taken to be the minimum of the data)

References

Bhatti SH, Hussain S, Ahmad T, Aslam M, Aftab M, Raza MA (2018). "Efficient estimation of Pareto model: Some modified percentile estimators." PLoS ONE, 13(5), 1-15.

Examples

```
x <- generate_pareto(10000, 5, 2)
alpha_geometric_percentile(x)
```

alpha_hills

Estimating the Shape Parameter by Hill's Estimator

Description

This function uses the Hill's Estimator to estimate the shape parameter of a given set of data. (Nair et al. 2019; Pokorna 2016; Hill 1975) It is especially useful when the data is known not to follow an exact Pareto distribution but the tail of the data does. Thus, the specification of k, the kth largest observation, allows to specify the point from where Pareto-like behavior may be seen. It is also possible to specify the value at which the tail begins. When k=n, the Hill's Estimator returns the same estimate as `alpha_mle` with a warning notifying the user.

Usage

```
alpha_hills(dat, k, value = FALSE)
```

Arguments

<code>dat</code>	vector of observations
<code>k</code>	number of observations / value equal to or greater than to consider for tail
<code>value</code>	(TRUE/FALSE) indicating if the value which is specified in "k" (TRUE)

Value

A list of the following form:

shape Estimate of the shape parameter of the data

scale Estimate of the scale parameter of the data (which is taken to be the minimum of the data)

References

Nair J, Wierman A, Zwart B (2019). "The Fundamentals Of Heavy Tails: Properties, Emergence, And Identification." <http://users.cms.caltech.edu/adamw/heavytails.html>.

Pokorna M (2016). Estimation and Application of the Tail Index. Bachelor's thesis, Charles University in Prague, Faculty of Social Sciences, Institute of Economic Studies.

Hill B (1975). "A Simple General Approach To Inference About The Tail Of A Distribution." The Annals of Statistics, 3(5), 1163-1174.

Examples

```
x <- generate_pareto(10000, 5, 2)
alpha_hills(x, 400)
```

alpha_ls

Estimating the Shape Parameter by Method of Least Squares

Description

This function uses the Method of Least Squares to estimate the shape parameter of a given set of data. (Zaher et al. 2014; Nair et al. 2019)

Usage

```
alpha_ls(dat)
```

Arguments

dat	vector of observations
-----	------------------------

Value

A list of the following form:

shape Estimate of the shape parameter of the data

scale Estimate of the scale parameter of the data (which is taken to be the minimum of the data)

References

Zaher HM, El-Sheik AA, El-Magd NATA (2014). "Estimation of Pareto Parameters Using a Fuzzy Least-Squares Method and Other Known Techniques with a Comparison." British Journal of Mathematics & Computer Science, 4(14), 2067-2088.

Nair J, Wierman A, Zwart B (2019). "The Fundamentals Of Heavy Tails: Properties, Emergence, And Identification." <http://users.cms.caltech.edu/adamw/heavytails.html>.

Examples

```
x <- generate_pareto(10000, 5, 2)
alpha_ls(x)
```

alpha_mle*Estimating the Shape Parameter by Method of Maximum Likelihood (MLE)***Description**

This function can be used to estimate the shape parameter using the Maximum Likelihood Estimator method (Newman 2005). It can be used to obtain biased and unbiased estimates of the shape and scale parameters as well as the confidence interval for the shape parameter for the biased estimates.

Usage

```
alpha_mle(dat, biased = TRUE, significance = NULL)
```

Arguments

dat	vector of observations
biased	TRUE/FALSE to indicate biased or unbiased estimates
significance	level of significance

Value

A list of the following form:

- shape** Estimate of the shape parameter of the data
- lower_bound** Upper error bound of the estimate of shape
- upper_bound** Lower error bound of the estimate of shape
- scale** Estimate of the scale parameter of the data (which is taken to be the minimum of the data)

References

Newman MEJ (2005). "Power Laws, Pareto Distributions And Zipf's Law." *Contemporary Physics*, 46, 323-351.

Examples

```
x <- generate_pareto(10000, 5, 2)
alpha_mle(x, TRUE, 0.05)

x <- generate_pareto(10000, 5, 2)
alpha_mle(x, FALSE)
```

alpha_modified_percentile*Estimating the Shape Parameter by Method of Modified Percentiles***Description**

This function uses the Method of Modified Percentiles to estimate the shape parameter of a given set of data. (Bhatti et al. 2018)

Usage

```
alpha_modified_percentile(dat)
```

Arguments

dat	vector of observations
-----	------------------------

Value

A list of the following form:

shape Estimate of the shape parameter of the data

scale Estimate of the scale parameter of the data (which is taken to be the minimum of the data)

References

Bhatti SH, Hussain S, Ahmad T, Aslam M, Aftab M, Raza MA (2018). "Efficient estimation of Pareto model: Some modified percentile estimators." PLoS ONE, 13(5), 1-15.

Examples

```
x <- generate_pareto(10000, 5, 2)
alpha_modified_percentile(x)
```

alpha_moment*Estimating the Shape Parameter by Method of Moments***Description**

This function uses the Method of Moments to estimate the shape parameter of a given set of data. (Rytgaard 1990) The method of moments is only accurate if α (shape parameter) is greater than or equal to 1 (Brazauskas and Serfling 2000). This function issues a warning if it detects that α may be less than 1.

Usage

```
alpha_moment(dat)
```

Arguments

dat vector of observations

Value

A list of the following form:

shape Estimate of the shape parameter of the data

scale Estimate of the scale parameter of the data (which is taken to be the minimum of the data)

References

Rytgaard M (1990). "Estimation In The Pareto Distribution." ASTIN Bulletin: The Journal Of The IAA, 20(2), 201-216.

Brazauskas V, Serfling R (2000). "Robust and Efficient Estimation Of The Tail Index Of A Single-Parameter Pareto Distribution." North American Actuarial Journal, 4, 12-27.

Examples

```
x <- generate_pareto(10000, 5, 2)
alpha_moment(x)
```

alpha_percentile

Estimating the Shape Parameter by Method of Percentiles

Description

This function uses the Method of Percentiles to estimate the shape parameter of a given set of data. (Bhatti et al. 2018)

Usage

```
alpha_percentile(dat)
```

Arguments

dat vector of observations

Value

A list of the following form:

shape Estimate of the shape parameter of the data

scale Estimate of the scale parameter of the data (which is taken to be the minimum of the data)

References

Bhatti SH, Hussain S, Ahmad T, Aslam M, Aftab M, Raza MA (2018). "Efficient estimation of Pareto model: Some modified percentile estimators." PLoS ONE, 13(5), 1-15.

Examples

```
x <- generate_pareto(10000, 5, 2)
alpha_percentile(x)
```

alpha_wls

Estimating the Shape Parameter by Weighted Least Squares Method (WLS)

Description

This function uses the Weighted Least Squares Method (WLS) to estimate the shape parameter of a given set of data. (Nair et al. 2019)

Usage

```
alpha_wls(dat)
```

Arguments

dat vector of observations

Value

A list of the following form:

shape Estimate of the shape parameter of the data

scale Estimate of the scale parameter of the data (which is taken to be the minimum of the data)

References

Nair J, Wierman A, Zwart B (2019). "The Fundamentals Of Heavy Tails: Properties, Emergence, And Identification." <http://users.cms.caltech.edu/adamw/heavytails.html>.

Examples

```
x <- generate_pareto(10000, 5, 2)
alpha_percentile(x)
```

generate_all_estimates*Obtain estimates for Parameters of Pareto Data from all methods*

Description

This function combines the results of all the methods (included in this package) provided to estimate the shape and scale parameters of the Pareto data and provides the results in a data frame. Hill's Estimator is not used in this comparison as it discards a set of observations. We also note here that when considering the entire data set, Hill's Estimate is equivalent to the MLE.

Usage

```
generate_all_estimates(dat)
```

Arguments

dat	vector of observations
-----	------------------------

Value

Dataframe with the following columns:

Method.of.Estimation Name of the method used for estimation

Shape.Parameter Estimates of the shape parameter of the data

Scale.Parameter Estimates of the scale parameter of the data

Examples

```
x <- generate_pareto(10000, 5, 2)
generate_all_estimates(x)
```

generate_pareto*Generating data from a Pareto Distribution.*

Description

This function is able to generate random Pareto distributed data with the specified shape and scale parameters. The function has been written to be similar in type to the popular runif and rexp type of functions for generating data from a particular distribution.

Usage

```
generate_pareto(sample_size, shape, scale)
```

Arguments

sample_size	number of observations
shape	shape parameter
scale	scale parameter

Value

Vector of Pareto distributed data of sample size `sample_size` with shape parameter `shape` and scale parameter `scale`.

Examples

```
generate_pareto(10000, 5, 2)
generate_pareto(100, 15, 6)
```

pareto_qq_test

*Q-Q Plot to test for Pareto Distribution***Description**

This function can be used as a first step to identify whether the data is Pareto distributed before estimating the tail index. If most of the data points appear to be distributed along a line, it is possible that the data may be Pareto. Conversely, if most of the data are distributed non-linearly, then the data is most probably not Pareto distributed.

Usage

```
pareto_qq_test(dat)
```

Arguments

dat	Data to be tested for Pareto distribution
-----	---

Details

This function plots the quantiles of the standard exponential distribution on the x-axis and the log values of the provided data on the y-axis. If Pareto data was supplied, a log transformation of this data would result in an exponential distribution with mean α . These data points would then show up on the QQ-plot as a line with slope $1/\alpha$.

The function makes use of the `plotly` package if available and installed or if not, defaults to the standard R plot.

Value

A Q-Q plot either using `plotly` if package is available or else a standard R plot.

Examples

```
x <- generate_pareto(10000, 5, 2)
pareto_qq_test(x)
```

pareto_test

Goodness of Fit Test for Pareto Distribution

Description

The *pareto_test* function can be used to identify whether the data is Pareto distributed (Gulati and Shapiro 2008). The test generates a p-value corresponding to the actual distribution of the data and is tested for significance. In the case of Pareto data, the p-value should be greater than the pre-determined significance level (generally taken as 0.05).

Usage

```
pareto_test(dat)
```

Arguments

dat	vector of observations
-----	------------------------

Value

A list of the following form:

p-value p-value indicating significance of the test

References

Gulati S, Shapiro S (2008). "Goodness-of-Fit Tests for Pareto Distribution." In F Vonta (ed.), Statistical Models and Methods for Biomedical and Technical Systems, chapter 19, pp. 259-274. Birkhauser Basel. ISBN 978-0-8176-4619-6. doi:10.1007/978-0-8176-4619-6.

Examples

```
x <- generate_pareto(10000, 5, 2)
pareto_test(x)
```

Index

alpha_geometric_percentile, 2, 3
alpha_hills, 2, 4
alpha_ls, 2, 5
alpha_mle, 2, 6
alpha_modified_percentile, 2, 7
alpha_moment, 2, 7
alpha_percentile, 2, 8
alpha_wls, 2, 9

generate_all_estimates, 2, 10
generate_pareto, 2, 10

pareto_qq_test, 2, 11
pareto_test, 2, 12
ptsuite-package, 2